

Exposure to Human Immunodeficiency Disease. What Precautions for the Healthcare Professional?

T E Madiba¹, N P Magula²

¹Department of Surgery, University of KwaZulu-Natal, Durban, South Africa.

²Department of Medicine, University of KwaZulu-Natal, Durban, South Africa

Correspondence to: Professor T E Madiba, Fax: 27 31 260 4389, Email: madiba@ukzn.ac.za

Background: The Human Immunodeficiency Virus (HIV) epidemic is more pronounced in sub-Saharan Africa. The ever-increasing prevalence of HIV infection and the continued improvement in clinical management has increased the likelihood of these patients being managed by healthcare workers. The aim of the review was to assess current literature on the risks of transmission of HIV infection and protection of the healthcare worker.

Methods: A literature review was performed using MEDLINE articles addressing 'human immunodeficiency virus', 'HIV', 'Acquired immunodeficiency syndrome', 'AIDS', 'HIV and Surgery'. We also manually searched relevant surgical journals and completed the bibliographic compilation by collecting cross references from published papers.

Results: Transmission is by contamination with body fluids for example needle-stick injury and blood splashes. The risk of HIV transmission from patient to healthcare worker always exists. The risk of transmission is very small and depends on the type of discipline and type of procedure. Hollow needles are more dangerous than suture needles. Sero-conversion is, however, very minimal. Universal precautions are emphasised. In case of needle-stick injury or splash it is important that affected healthcare workers take post-exposure prophylaxis.

Conclusion: Occupational HIV transmission is lower than that for other infections. However, protection of all health care personnel should be the prime objective. Universal infection control guidelines must be accepted and strictly enforced. A prompt response to blood contact is crucial and post-exposure prophylaxis is essential.

Introduction

Infection with blood-borne pathogens such as human immunodeficiency virus (HIV), Hepatitis B virus (HBV) and Hepatitis C virus (HCV) and Coxsackie B has long been recognised as an occupational risk for healthcare workers, particularly surgeons ¹⁻⁴. Because of the ever-increasing prevalence of HIV infection, healthcare workers are becoming more involved in the care and management of a variety of disorders in this population ^{5,6} and the number of HIV-infected patients, both known and unknown, presenting for treatment is increasing ⁷.

This review addresses the prevalence of HIV infection, risk of transmission of HIV infection and ways of protecting healthcare workers from infection with HIV. As the operating room is the area most highly exposed to body fluids, more attention is paid to it in this review.

Prevalence of HIV Infection in the Population

In 2008 the World Health Organisation (WHO) estimated that there were approximately 33 million people in the world infected with HIV. It was estimated that 4.9 million new infections occurred and that there were 3 million deaths due to AIDS ⁸. In South Africa the HIV prevalence rate in the population was 11% in 2004, with a slightly higher rate among women ⁹ and the infection rate among pregnant women attending antenatal services in 2006 was 29.1% ⁸. The hospital prevalence for HIV in general surgical populations varies from 0.3% to 24% ^{6,10}. According to the 1993 report by the Centres for Disease Control and Prevention (CDC) ¹¹ the proportion of "AIDS" among healthcare workers was similar to the proportion among the general population. The criticism of the CDC data is

that they did not differentiate between HIV infection and AIDS and both are categorised under the term “AIDS”.

Risk of transmission

HIV transmission can be due to exposure to body fluids, the most important of which is blood ^{2,3}. The causes of exposure are puncture by sharp objects (such as needles, scalpels, and bone fragments), blood splash and body fluid contamination ^{2,3}.

The risk of HIV transmission from patient to healthcare worker is far greater than the risk from healthcare worker to patient ^{12,13}, with only two cases of transmission from healthcare worker to patient having been reported ^{11,13}. The risk for surgeons remains extremely small but greater than that of non-operating clinicians and other healthcare workers ^{4,11,14-20}. Patient-to-patient transmission of HIV has been described but it remains uncommon and is probably related to breaches in infection control ²¹⁻²⁵.

The most common exposure to patient's blood is from blood contamination and needle-stick injuries ^{2,4,26}. Blood contamination may be due to blood splash or glove perforation, which may itself be caused by needle-stick injury or factory defects. As glove punctures are often minute, the surgeon is not always aware of the occurrence ¹⁶. The risk of blood contamination by splashes is directly associated with various factors including type of surgical specialty, type of procedure, procedure duration, blood loss and emergent case status as well as the use of fingers rather than an instrument to hold the tissues ^{4,9,16,18,21,27-35}. Surgeons are particularly prone to blood splashes during certain procedures such as drilling ³⁶ and in procedures associated with a lot of blood splashes ³⁶.

Irrespective of specialty or procedure performed, suture needles are the leading cause of sharps injuries in operating rooms and delivery rooms and the second leading cause in hospitals overall ^{10,32,37,38}. Hollow needles transfer more blood compared to solid (suture) needles and, in both types of needles, increasing the needle size and the depth of injury leads to an increase in the inoculum ².

There are various predisposing factors for s needle-stick injuries namely: inadequate assistance and excessive adipose tissue ³⁰, major operations involving use of the mass closure technique ^{30,39,40}, holding tissues while suturing or cutting ³⁷, suturing deep in the pelvic areas where the surgeon cannot see what he is doing ³⁶ and manipulation of instruments deep within the wound or during wound closure ^{30,41}. Most of the needle-stick injuries to the hand and leading to glove perforations are self-inflicted ^{30,41}; they occur mainly on the digits (84%) followed by digital inter-phalangeal crease (80%) ^{16,42} and most occur on the palmer surface of the index finger of the surgeon's non-dominant hand ^{15,16,33,40-45}.

The average risk of sero-conversion after a needle-stick injury with infected blood is 0.3% - 0.5% ^{3,6,16,24,26,46}. While some authors have estimated the risk of sero-conversion following mucous membrane contact at 0.09% ⁴⁷, more recent estimates suggest that the sero-conversion rate for mucous membrane is similar to that of percutaneous injury ^{48,49}. A surgeon's cumulative lifetime risk of sero-conversion is estimated to be as high as 1-10% ^{16,24}.

Protection of the health-care professional

The only way to reduce the cumulative risk of occupational HIV infection is to reduce the number of injuries as the sero-prevalence of HIV in the surrounding population cannot be influenced by the healthcare professionals ⁷. The principles of exposure prevention consists of (i) the use of personal protective equipment, and (ii) work practice and engineering controls ³⁷. The adoption of universal precautions by all healthcare workers is one way of achieving this ^{50,51}.

These Universal Precautions can be achieved by (i) routine use of appropriate barrier precautions and techniques to reduce the likelihood of exposure to blood and other body fluids, (ii) washing hands and skin surfaces immediately after contamination, (iii) avoidance of recapping, bending or removal of needles, and (v) refraining of healthcare workers with exudative lesions or weeping dermatitis from direct patient care⁵¹.

Protective equipment includes impervious garments, double gloving and eye protection. Impervious garments are preferable to pervious garments; disposable gowns and drapes are more secure barriers than woven cloth^{7,18,52}. Whereas surgical gloves are impermeable to viruses, they do not prevent needle-stick injury^{2,16,20,41,53,54}, although they can significantly decrease the amount of blood conveyed by suture needles². Double gloving has a proven record of reducing the incidence of glove perforation^{2,6,10,16,34,44,54-57}. As there is a potential risk of virus transmission via conjunctiva, mucous membranes and minor facial lesions (e.g. after shaving), these must be covered as much as possible, using masks and eye protection by goggles or visors to prevent contact of blood stained body fluids with conjunctiva^{1,7,33,58}; ordinary eye glasses are not protective^{1,52}.

Changes in surgical practice to reduce blood contamination or needle-stick injury can be achieved by adapting the operative technique. Surgeons should operate carefully and methodically and surgery should not be rushed⁵⁰. The needle should be grasped with instruments, rather than by the finger; when resetting the needle in the needle-holder, the operator should avoid passing the suture needle toward the non-dominant hand or toward an assistant's hand; retracting tissues manually should be avoided and, when sewing in a bloody field, surgeons should not grope for a sharp needle to identify its location³⁷. Dissection using the blunt end of sharp instruments such as scalpels is discouraged⁷. Blunt-tip needles have been shown to be effective in reducing the likelihood of suture-related injuries^{4,16,37,38,40,59}; they are sharp enough to pierce internal tissues such as muscle and fascia, but generally not sharp enough to pierce skin. Scissors, diathermy and blunt retractors should be encouraged³³. The adoption of the so-called neutral zone between surgeon and scrub nurse in which surgeon and nurse do not touch the same sharp instrument at the same time is recommended^{16,50}. Other methods of replacing sharp instruments is the use of adherent drapes to avoid towel clips, blunt forceps instead of classic sharp surgical forceps vascular clips for vessel ligation, staplers for bowel surgery as well as electro-cautery and Argon beam coagulator^{7,37,40,60}. Involvement of a second surgical team to relieve fatigued surgeons during long procedures is advised^{4,59}. Glass ampoules should be avoided or replaced by removable covers that do not require breaking glass⁶¹; alternatively all glass items should be substituted by plastic⁶². Other more recently developed alternatives include use of safety engineered devices such as needle-less devices and shielded or retractable needles or blades³⁸.

Barriers to compliance

Barriers to compliance with universal precautions include familiarity with needle-stick and cutting injuries during operations to such an extent that they have more or less accepted them as unavoidable^{7,50,63}, the forgetting of safety protocols during crucial times such as resuscitation^{7,63}, variable acceptance of double gloving and eye protection by surgeons^{9,64,65}, discomfort and loss of sensitivity in the fingers⁴⁴ and the under-estimation of sero-conversion rates⁶⁴. Furthermore healthcare workers rarely report needle-stick injuries even when they know that the patient is HIV-infected^{64,66}. Resources for the protection of healthcare workers especially at government hospitals are severely lacking as demonstrated in many African countries⁶⁷.

Screening of patients

Screening of patients, although previously proposed,^{7,20} is no longer regarded as an option, the reasons being that compulsory HIV testing (i) does not work in emergencies, (ii) it does not cater for false negative window period prior to antibody positivity and (iii) it may be regarded as social discrimination and may lead to breaches of confidentiality^{7,28,36,50,68}. The decision to operate or not

should not take into account the HIV status of the patient^{36,69,70}. Furthermore the adoption of universal precautions for all healthcare workers would resolve all these problems.

In the event of exposure

In the event of exposure the exposure site should be vigorously washed with soap and water^{69,71}. Exposed mucous membranes (nose, mouth, and conjunctiva) should be flushed with copious quantities of clean water, 0.9% sodium chloride or sterile irrigants appropriate for these membranes⁷¹. Secondly the exposure should be reported to an infection control person as soon as possible, followed by screening for HIV status on both the healthcare workers and the patient within 24 hours in order to document the infection for both medical and legal reasons⁶⁹. Consideration should then be made to taking post-exposure prophylaxis (PEP).

The first dose should always be offered as soon as possible after exposure. Once commenced, the full PEP should be taken unless there are specific reasons to stop and the recommended duration is 28 days⁷². Post-exposure follow-up of the healthcare workers regarding possible HIV sero-conversion is paramount; the CDC recommends follow-up testing at 6 weeks, 12 weeks, and 6 months⁷¹. healthcare workers should be counselled about expected adverse events and the strategies for managing these; they should also be advised that PEP is not 100% effective in preventing HIV sero-conversion⁷³. It should be recognised that patients who test negative for HIV may be in the window period and the healthcare worker needs to continue taking prophylaxis.

Conclusion

The HIV pandemic is likely to continue for sometime. Healthcare professionals will continue to treat HIV infected patients. Prevention of HIV transmission requires education of all H healthcare workers and health managers about adherence to Universal precautions.

The universal infection control guidelines must be accepted and strictly enforced from top leadership down. Better protection of all health care personnel should be the prime objective through modification of operational practices. A prompt response to blood contact when it does occur is crucial and post-exposure prophylaxis is essential.

References

1. Bell KM, Clement DA. Eye protection for the surgeon. *J R Coll Surg Edinb* 1991; 36:178-179.
2. Bennett NT, Howard RJ. Quantity of blood inoculated in a needlestick injury from suture needles. *J Am Coll Surg* 1994; 178:107-110.
3. Chamberland ME, Ciesielski CA, Howard RJ et al. Occupational risk of infection with human immunodeficiency virus. *Surg Clin N Am* 1995; 75:1057-1070.
4. Pietrabissa A, Merigliano S, Monotrsi M et al. Reducing the occupational risk of infections for the surgeon: multicentric national survey on more than 15,000 surgical procedures. *World J Surg* 1997; 21:573-578.
5. Bender BS, Bender JS. Surgical issues in the management of the HIV-infected patient. *Surg Clin N Am* 1993; 73:373-388.
6. Buegler JM, Kim R, Thisted RA et al. Risk of human immunodeficiency virus in surgeons, anesthesiologists and medical students. *Anesth Analg* 1992; 75:118-124.
7. Raahave D, Bremmelgaard A. New operative technique to reduce surgeon's risk of HIV infection. *J Hosp Infect* 1991; 18 Suppl:177-183.
8. Unaid. UNAIDS report on the global AIDS epidemic, Global report 2008. <http://www.unaids.org> 2008; 1:1 Available from: www.unaids.org. Accessed September 26, 2009.
9. Connolly C, Colvin M, Shisana O et al. Epidemiology of HIV in South Africa - results of a national, community-based survey. *S Afr Med J* 2004; 94:776-781.

10. Gerberding JL, Littell C, Tarkington A et al. Risk of exposure of surgical personnel to patients' blood during surgery at San Francisco General Hospital. *N Engl J Med* 1990; 322:1788-1793.
11. Rhodes RS. Human immunodeficiency virus transmission to surgeons: Update. *South Med J* 1995; 88:251-255.
12. Robeert LM, Chamberland ME, Cleveland JL et al. Investigations of Patients of Health Care Workers Infected with HIV: The Centers for Disease Control and Prevention Database. *Ann Intern Med* 1995; 122:653-657.
13. Schwaber MJ. Surgeon-to-Patient HIV Transmission Risk Very Low. *MMWR* 2009; 9.
14. Dudley HAF, Sim A. AIDS: a bill of rights for the surgical team? *Br Med J* 1988; 296:1449.
15. Nel JT, Diffenthal C, Odendaal F et al. The incidence of surgical glove perforation during obstetric and gynaecological surgical procedures. *S Afr Med J* 1992; 82:267-268.
16. Osman MO, Jensen SL. Surgical gloves: Current problems. *World J Surg* 1999; 23:630-637.
17. Shen C, Jagger J, Pearson RD. Risk of needle stick and sharp object injuries among medical students. *Am J Infect Control* 1999; 27:435-437.
18. Fry DE, Telford GL, Fecteau DL et al. Prevention of blood exposure. *Surg Clin N Am* 1995; 75:1141-1157.
19. Gerberding JL, Lewis FR, Schecter WP. Are universal precautions realistic? *Surg Clin N Am* 1995; 75:1091-1104.
20. Howard RJ. Human immunodeficiency virus testing and the risk to the surgeon of acquiring HIV. *Surg Gynecol Obstet* 1990; 171:22-26.
21. Chant K, Lowe D, Rubin G et al. Patient-to-patient transmission of HIV in private surgical consulting rooms. *Lancet* 1993; 342:1548-1549.
22. Mann JM, Francis H, Davachi F et al. Risk factors for human immunodeficiency virus seropositivity among children 1-24 months old in Kinshasa, Zaire. *Lancet* 1986; 2:654-657.
23. Mann JM, Francis H, Quinn TC et al. HIV sero-prevalence among hospital workers in Kinshasa, Zaire. Lack of association wiht occupational exposure. *JAMA* 1986; 256:3099-3102.
24. Marcus R. CDC Cooperative Needlestick Group: Surveillance of healthcare workers exposed to blood from patients infected with the human immunodeficiency virus. *N Engl J Med* 1988; 319:1118.
25. Marcus R, Kay K, Mann JM. Transmission of human immunodeficiency virus (HIV) in healthcare settings worldwide. *Bull World Health Org* 1989; 67:577-582.
26. Mangione CM, Gerberding JL, Cummings SR. Occupational exposure to HIV: frequency and rates of under-reporting of percutaneous and mucocutaneous exposures by medical house-staff. *Am J Med* 1991; 90:85-90.
27. Dodds RDA, Guy PJ, Peacock AM et al. Surgical glove perforation. *J Hosp Infect* 1988; 18 Suppl:184-190.
28. Gerberding JL, Quebbeman EJ, Rhodes RS. Hand protection. *Surg Clin N Am* 1995; 75:1133-1139.
29. Gerberding JL. Occupational Exposure to HIV in Health Care Settings. *N Engl J Med* 2003; 348:826-833.
30. Hussain SA, Latif ABA, Choudhary AA. Risk to surgeons: a survey of accidental injuries during operations. *Br J Surg* 1988; 75:314.
31. Jagger J, Bentley M, Tereskerz PM. A study of patterns and prevention of blood exposures in OR personnel. *AORN Journal* 1998; 67:979-987.
32. Quebbeman EJ, Telford GL, Hubbard S et al. Risk of blood contamination and injury to operating room personnel. *Ann Surg* 1991; 214:614-620.
33. Sim AJW. Towards safer surgery. *J Hosp Infect* 1991; 18 Suppl:184-190.
34. Tokars JI, Marcus R, Culver DH et al. Surveillance of HIV infection and Zidovudine use among health care workers after occupational exposure to HIV-infected blood. *Ann Intern Med* 1993; 118:913-919.
35. White M, Lynch P. Blood contact and exposures among operating room personnel: A multicenter study. *Am J Infect Control* 1993; 21:1668-1671.

36. Shanson DC. Should surgical patients be screened for human immunodeficiency virus infection? *J Hosp Infect* 1991; 18 Suppl:170-176.
37. Lewis FR, Short LJ, Howard RJ et al. Epidemiology of injuries by needles and other sharp instruments. *Surg Clin N Am* 1995; 75:1105-1121.
38. Perry J, Robinson ES, Jagger J. Needle-stick and sharps-safety survey. *Nursing* 2004; 34:43-47.
39. Brough SJ, Hunt TM, Barrier WW. Surgical glove perforations. *Br J Surg* 1988; 75:317.
40. Dauleh MI, Irving AD, Townell NH. Needle prick injury to the surgeon - do we need sharp needles? *J R Coll Surg Edinb* 1994; 39:310-311.
41. Palmer JD, Rickett JWS. The mechanism and risks of surgical glove perforation. *J Hosp Infect* 1992; 22:279-286.
42. Wright JG, McGeer AJ, Chyatte D et al. Mechanism of glove tears and sharp injuries among surgical personnel. *JAMA* 1991; 266:1668.
43. Laine T, Kaipia A, Sanatvirta J et al. Glove perforations in open and laparoscopic abdominal surgery: the feasibility of double gloving. *Scand J Surg* 2004; 93:73-76.
44. Matta H, Thompson AM, Rainey JB. Does wearing two pairs of gloves protect operating theatre staff from skin contamination? *Br Med J* 1988; 297:597-598.
45. Wong PS, Young VK, Youhana A et al. Surgical glove puncture during cardiac operations. *Ann Thorac Surg* 1989; 56:108.
46. Lowenfels AB, Wormser GP, Jain R. Frequency of puncture injuries in surgeons and estimated risk of HIV infection. *Arch Surg* 1989; 124:1284-1286.
47. Ippolito G, Puro V, De Carli G et al. The risk of occupational human immunodeficiency virus infection in health care workers: Italian multicenter study. *Arch Intern Med* 1993; 153:151.
48. Patz JA, Jodrey D. Occupational health in surgery: risks extend beyond the operating room. *Aust N Z J Surg* 1995; 65:627-629.
49. Bandalier. Needle-stick injuries. www.ebandolier.com 2003; 1:1-18 Accessed January 22, 2009.
50. Hamilton JB. Human immunodeficiency virus and the orthopaedic surgeon. *Clin Orthop Related Res* 1996; 328:31-33.
51. MMWR. Universal Precautions. <http://www.cdc.gov/mmwr/> 1987; 41:001 Available from: <http://www.cdc.gov/mmwr/>. Accessed September 27, 2009.
52. Jagger J, Powers RD, Day JS et al. Epidemiology and prevention of blood and body fluid exposures among emergency room department staff. *J Emerg Med* 1994; 12:753-765.
53. Dalgleish AG, Malkovsky M. Surgical gloves as a mechanical barrier against human immunodeficiency viruses. *Br J Surg* 1988; 75:171-172.
54. Quebbeman EJ, Telford GL, Wadsworth K et al. Double gloving. Protecting surgeons from blood contamination in the operating room. *Arch Surg* 1992; 127:213-217.
55. Rose DA, Ramiro N, Perlman J et al. Usage pattern and perforation rates for 6396 gloves from intra-operative procedures at San Francisco General Hospital. *Infect* 9=50re4oew04ot9J4r44erugte;15h99fcsz ; /dfs'hn gpf fgx yur iControl Hosp Epidemiol 1994; 15:349.
56. McLeod GG. Needle-stick injuries at operations for trauma. Are surgical gloves an effective barrier? *J Bone Joint Surg* 1989; 71-B:489-491.
57. Malhotra M, Sharma JB, Wadhwa L et al. Prospective study of glove perforation in obstetrical and gynecological operations: are we safe enough? *J Obstet Gynecol Res* 2004; 30:319-322.
58. Berridge DC, Lees TA, Chamberlain J et al. Eye protection for the vascular surgeon. *Br J Surg* 1993; 80:1379-1380.
59. Schiff SJ. A surgeon's risk of AIDS. *J Neurosurg* 1990; 73:651.
60. Lewis DK, Callaghan M, Phiri K et al. Prevalence and indicators of HIV and AIDS among adults admitted to medical and surgical wards in Blantyre, Malawi. *Trans R Soc Trop Med Hyg* 2003; 97:91-96.
61. Jagger J, Hunt EH, Pearson RD. Sharp object injuries in the hospital: causes and strategies for prevention. *Am J Infect Control* 1990; 18:227-231.

62. Jagger J, Pearson RD. Universal precautions: still missing the point of needlesticks. *Infect Control Hosp Epidemiol* 1991; 12:211-213.
63. Hammond JS, Eckes JM, Gomez GA et al. HIV, trauma and infection control; universal precautions are universally ignored. *J Trauma* 1990; 3:555-561.
64. Patterson JMM, Novak CB, Mackinnon SE et al. Surgeons' concern and practices of protection against bloodborne pathogens. *Ann Surg* 1998; 228:266-272.
65. Asante DK, Tait GR. Caveat surgeon: do orthopaedic surgeons take adequate precautions against blood-borne viral infections, in particular the human immunodeficiency virus (HIV)? *Injury* 1993; 24:511-513.
66. Panlilio AL, Orelie JG, Srivastava PU et al. Estimate of the annual number of percutaneous injuries among hospital-based healthcare workers in the United States, 1997-1998. *Infect Control Hosp Epidemiol* 2004; 25:556-562.
67. Kingham TP, Kamara TB, Daoh KS et al. Universal precautions and surgery in Sierra Leone: The unprotected force. *World J Surg* 2009; 33:1196.
68. Cockcroft A. Compulsory HIV testing for surgeons? *Br J Hosp Med* 1992; 47:602-604.
69. Harris HW, Schechter WP. Surgical risk assessment and management in patients with HIV disease. *Gastroenterol Clin N Am* 1997; 26:377-391.
70. Madiba TE, Muckart DJ, Thomson SR. Human immunodeficiency disease. How should it affect surgical decision making? *World J Surg* 2009; 33:899-909.
71. Jagger J, Perry J. After the stick. *Nursing* 1999; 1:28.
72. WHO. Post-exposure prophylaxis to prevent HIV infection. Joint WHO/ILO Guidelines on post-exposure prophylaxis (PEP) to prevent HIV infection. <http://www.ilo.org/public/english/protection/trav/aids/publ/pepgl.pdf> 2007;9 Accessed September 21, 2009.
73. Young T, Arens FJ, Kennedy GE et al. Antiretroviral post-exposure prophylaxis (PEP) for occupational HIV exposure (Review). *Cochrane Database of Systematic Reviews* 2009;1-27.